arXiv:physics/0110009v2 [physics.geo-ph] 28 May 2002

On the Doppler distortion of the sea-wave spectra

A. O. Korotkevich

Landau Institute for Theoretical Physics, Moscow 117334, Kosygin Street 2, Russia

Manuscript submitted to
Nonlinear Processes in Geophysics
February 2, 2008

Offset requests to:
A. O. Korotkevich, kao@landau.ac.ru

Abstract

Discussions on a form of a frequency spectrum of wind-driven sea waves just above the spectral maximum continue during the last three decades. In 1958 Phillips made a conjecture that wave breaking is the main mechanism responsible for the spectrum formation. That leads to the spectrum decay $\sim \omega^{-5}$, where ω is the frequency of the waves. There is a contradiction between the numerous experimental data and this spectrum. The experiments show decay $\sim \omega^{-4}$. There are two general ways of the explanation of this phenomenon. The first one (proposed by Banner (1990)) takes into account the Doppler effect due to surface circular currents generated by long waves in the Phillips model. The second approach ascends to the work by Zakharov and Filonenko (1968). It is based on four-wave interactions in the kinetic equation and gives good agreement with the experimental data.

In this article the contribution to the Phillips model due to the Doppler effect was examined. We considered the cases of isotropic and strongly anisotropic spectra. Both analytical results and numerical computations give us an evidence to state that the Doppler effect cannot explain the divergence of Phillips model with experimental data.

Correspondence to: A. O. Korotkevich, kao@landau.ac.ru

1 Introduction

There is a long discussion in the scientific world about the form of wind generated sea wave spectrum in the ”universal” region of frequencies $\omega \gtrsim \omega_p$, where ω_p is the frequency of the spectral maximum.

There are two main points of view. The first one proposed by Phillips in 1958 gives us the Phillips' spectrum in universal region

$$F(\omega) \simeq \frac{\alpha g^2}{\omega^5}, \tag{1}$$

where α is dimensionless constant, which according to experimental data appears to be rather small $\alpha \simeq 0.01$. The second point of view introduced by Zakharov and Filonenko (1966) gives us in universal region following spectrum

$$F(\omega) \simeq \frac{\beta \epsilon^{\frac{1}{2}} g v}{\omega^4}, \tag{2}$$

where β - is dimensionless constant, v - is wind velocity, $\epsilon \simeq \frac{\rho_a}{\rho_w}$ - is the ration between ocean water and atmosphere densities.

The (1) and (2) formulas are based upon completely opposite propositions. The Phillips' spectrum (1) takes place if the spectrum at the high frequency region is mainly determined by appearance of sharp crests and wave breaking. The level of nonlinearity depending on steepness $\mu = ka$ in this case at least at the level of the wave peaks is extremely high - $\mu \simeq 0.44$.

On the other side spectrum (2) is based on presumption of small mean steepness - $\mu \ll 1$. This proposition is in great concordance with observed data. Usually $\mu \lesssim 0.1$.

Following this ideology the wind generated waves' ensemble is described by Hasselman kinetic equation (Hasselman 1962,1963)

$$\frac{\partial N}{\partial t} = S_{nl} + p^+ + p^-. \tag{3}$$

Zakharov-Filonenko spectrum appears as exact solution of the particular case of Eq. (3)

$$S_{nl} = 0, \tag{4}$$

this is a classical case of Kolmogorov-Zakharov (KZ) spectra. This weakly turbulent theory is far advanced both analytically and numerically (Zakharov and Filonenko 1967, 1966; Craik 1987; Zakharov at al. 1992; Pushkarev and Zakharov 1996, 2000).

A lot of experimental observations (Forristall, 1981; Kahma, 1981; Donelan, 1985, 1998, Hansen at al. 1990) beginning from famous work of Toba (1973) gives us evidences that in most cases at moderately high frequencies $\frac{\omega}{\omega_p} \lesssim 10$ the Zakharov-Filonenko spectrum (2) is well detected. Sometimes in the region of extremely high frequencies $\frac{\omega}{\omega_p} \gg 10$ the Phillips spectrum (1) takes place.

”Universality” of spectrum ω^{-4} became obvious in the middle of eighties. In 1986 O. Phillips, ”father” of ω^{-5} spectrum published an article with confession that spectrum (1) may takes place only at high frequency region and is not universal.

In the region of moderate frequencies the ω^{-4} spectrum is dominating with confidence. From this fact the question of physical interpretation of spectrum $F(\omega) \simeq \omega^{-4}$ becomes very important. The origin of this spectrum is rather obvious from our point of view. This spectrum is just weakly turbulent KZ-spectrum. There are another explanations, though.

One of them was proposed by Phillips in 1986. He made supposition that spectrum ω^{-4} is a result of balance of all three terms of Eq.(3)

$$S_{nl} \simeq p^{+} + p^{-}. \tag{5}$$

This affirmation unlikely can be treated as a serious theory, because the expression of p^{-} is completely unknown. There are only very rough empirical formulas. However in that paper this expression was taken in the form giving us the balance in Eq. (5). But there is no any ground for this assumption.

Another explanation was proposed by Banner (1990) and developed by Donelan (1998). They consider the ω^{-4} spectrum as an artifact of the observation of time histories at a point brought about by Doppler shifting of the short waves riding on the orbital currents of the long waves. Let us suppose that spatial spectrum of the short waves $\Phi(k)$ is known. The problem is to derive the frequency spectrum $F(\omega)$ using only $\Phi(k)$. Normally we have to assume that waves obey linear dispersion equation. In this case the following simple expression takes place

$$F(\omega) = \int \Phi(k)\delta(\omega - \omega(k))dk.$$

However in the presence of long waves the dispersion law of the short waves modifies due to Doppler effect at the orbital velocity's field $\mathbf{v}$ of the long wave component

$$\omega(k) \to \omega(k) + (\mathbf{kv}),$$

where $\mathbf{v}$ is the chaotic quasi-periodic function of time with the zero mean value. In this case the frequency spectrum $F(\omega)$ becomes wider.

The present paper is devoted to the quantitative theory of this without any doubt very important phenomenon. We consider it in a very simplified way, supposing that $\mathbf{v}$ is sinusoidal function of time, i.e. assuming that the long wave is strictly monochromatic. This supposition is not too unreal. Undoubtedly this gives us the correct order of magnitude. All the more the result of this paper is quite clearly understandable.

In Sect. 2 we introduced the general notations (following Phillips). In Sect. 3 the general formula for spectrum (taking into account Doppler shifting) is derived. It is convenient to introduce a new set of dimensionless variables, this is considered in Sect. 4. In Sect. 5 we develop an important case of dispersion $\omega^2 = gk$ ("deep water") where g is the gravity acceleration, k is the wave number equal to $2\pi/\lambda$. Section 6 is devoted to numerical results. The conclusion is placed in Sect. 7.

At the case of sufficiently small steepness $\mu \simeq 0.05$, the Doppler effect appears to be too weak to explain the clearly observed difference of ω^{-5} spectrum and experimental results, at least in the region of universal frequencies $\omega/\omega_p \simeq 5 - 10$. This fact makes too detailed inspection redundant.

However, it is worth to say, that at higher frequencies the Doppler shifting can play very important role. This fact cannot influence on interpretation of experimental data at moderate frequencies though. At our standpoint the only conclusive interpretation of observed spectrum ω^{-4} is treating it as weak turbulent Kolmogorov-Zakharov spectrum. It is worth to add that many direct measurements of spatial spectrum $\Phi(k) =< |\xi_k|^2 >$ by photographing of the ocean surface and completely excluding the Doppler effect influence give us in the region of moderately high wave numbers the expectable result

$$\Phi(k) \simeq k^{7/2}.$$

This fact is in great accordance with frequency spectrum $F(\omega) \simeq \omega^{-4}$ at the absence of Doppler shifting.

2 Formulation of the problem and notations

Following the notations of Phillips, the wave spectrum for homogeneous stationary wave field can be introduced as follows

$$X(\mathbf{k},\omega) = (2\pi)^{-3} \iint\limits_{-\infty}^{+\infty} \rho(\mathbf{r},t) \exp[-i(\mathbf{k}r - \omega t)] d\mathbf{r} dt, \tag{6}$$

where $\rho(\mathbf{r},t) = \overline{\xi(\mathbf{x},t_0)\xi(\mathbf{x}+\mathbf{r},t_0+t)}$ is the covariance of the surface displacement $\xi(\mathbf{x},t)$, $\mathbf{r}$ is the spatial separation vector, t is the time separation, $\mathbf{k} = \overrightarrow{(k_1,k_2)} = \overrightarrow{(k,\theta)}$ is the wavenumber vector (the second representation is just the same in polar coordinates) and ω is the radian frequency. Obviously it can be rewritten in terms of Fourier transforms of corresponding functions

$$\langle \xi(\mathbf{k},\omega)\xi^*(\mathbf{k}',\omega')\rangle = X(\mathbf{k},\omega)\delta_{\mathbf{k}-\mathbf{k}'}\delta_{\omega-\omega'}, \tag{7}$$

$$X(-\mathbf{k},-\omega) = X(\mathbf{k},\omega).$$

It is convenient to introduce additional spectra which are the reduced forms of Eq. (6)

$$\Phi(\mathbf{k}) = 2\int\limits_{0}^{+\infty} X(\mathbf{k},\omega) d\omega, \tag{8}$$

$$F(\omega) = 2\int\limits_{0}^{2\pi}\int\limits_{0}^{+\infty} X(\mathbf{k},\omega) k dk d\theta. \tag{9}$$

In general case it is impossible to express $F(\omega)$ in terms of $\Phi(\mathbf{k})$. But in the case of small wave amplitude we can use the linearized equation

$$X(\mathbf{k},\omega) = \Phi(\mathbf{k})\left(\delta(\omega-\omega_k) + \delta(\omega+\omega_k)\right), \tag{10}$$

where $\omega_k = \omega(k)$ is the dispersion law and δ is the Dirac delta-function. Now one can rewrite Eq. (9) in the following form

$$F(\omega) = 2\int\limits_0^{2\pi}\int\limits_0^{+\infty} \Phi(k,\theta)\left(\delta(\omega-\omega_k)+\delta(\omega+\omega_k)\right)kdkd\theta. \tag{11}$$

Let us remember that $\omega_k > 0$

$$F(\omega) = 2\int\limits_0^{+\infty}\int\limits_0^{2\pi} \Phi(k,\theta)\delta(\omega-\omega_k)d\theta kdk. \tag{12}$$

Using the rules of integration of δ-function one can obtain

$$\begin{aligned} &F(\omega) = \frac{2k}{|\omega'(k)|}\int\limits_0^{2\pi}\Phi(k,\theta)d\theta, \\ &k = k(\omega), \end{aligned} \tag{13}$$

where $k(\omega)$ is the inverse function for $\omega(k)$.

In the deep water case $\omega = \sqrt{gk}$

$$k = \frac{\omega^2}{g}, \omega'(k) = \frac{1}{2}\sqrt{\frac{g}{k}} = \frac{1}{2}\frac{g}{\omega}$$

$$F(\omega) = \frac{4\omega^3}{g^2}\int\limits_0^{2\pi}\Phi\left(\frac{\omega^2}{g},\theta\right)d\theta. \tag{14}$$

For an isotropic spectrum ($\Phi(k,\theta) = \Phi(k)$)

$$F(\omega) = \frac{8\pi\omega^3}{g^2}\Phi\left(\frac{\omega^2}{g}\right). \tag{15}$$

For a strongly anisotropic spectrum ($\Phi(k,\theta) = \Phi(k)\delta(\theta-\theta_0)$)

$$F(\omega) = \frac{4\omega^3}{g^2}\Phi\left(\frac{\omega^2}{g}\right). \tag{16}$$

If the spectrum is determined by discontinuities of spatial derivative (wedges) caused by wave breaking according to Phillips (1958), we have

$$\xi \simeq \frac{1}{k^2}, \Phi(k) \simeq \frac{1}{k^4}. \tag{17}$$

In the isotropic case substitution Eq. (17) to Eq. (15) gives

$$F(\omega) = 8\pi\alpha\frac{g^2}{\omega^5}. \tag{18}$$

Similar calculation for Eq. (16) gives us the following result

$$F(\omega) = 4\alpha\frac{g^2}{\omega^5}. \tag{19}$$

Here α is the so-called Phillips' constant.

If the wedges appear on each wave crest, α should not be too small ($\alpha \simeq 0.1$). Any attempts to fit Eq. (18) or Eq. (19) to experimental facts give for α at least one order of magnitude less.

3 General formula for spectrum with the account for the Doppler effect

Let us assume that we study waves on the background of periodic current $\mathbf{v}(t)$ beneath the surface. Taking into account the Doppler effect in the dispersion equation one can get the following substitution

$$\omega(\mathbf{k}) \longrightarrow \omega(\mathbf{k}) + \mathbf{kv}(t).$$

Suppose that this current is described by a one-dimensional periodic function

$$v(t) = v \cos \omega_0 t.$$

This allows us to rewrite our substitution in the following form

$$\omega_k \longrightarrow \omega(k) + kv \cos(\omega_0 t) \cos \psi,$$

where ψ is the angle formed by $\mathbf{k}$ and $\mathbf{v}$ vectors. Without loss of generality one can take $\psi = \theta$. Using this expression one can write mean value of expression (12) as

$$F(\omega) = \frac{2}{T} \int\limits_0^T \int\limits_0^{+\infty} \int\limits_0^{2\pi} \Phi(k, \theta) \delta(\omega - \omega(k) - kv \cos(\omega_0 t) \cos \theta) d\theta k dk dt, \tag{20}$$

where T is the period equal to $2\pi/\omega_0$. It is useful to take the average with respect to t as the first step. Let us denote this factor to the part of the integral as M

$$M = \frac{1}{\pi} \int\limits_0^T \delta(\omega - kv \cos \omega_0 t \cos \theta - \omega_k) d(\omega_0 t).$$

The δ-function in this expression gives us the equation

$$\omega - \omega_k = kv \cos \theta \cos(\omega_0 t_0), \tag{21}$$

This equation have the following roots

$$\omega_0 t_0 = \pm \arccos \frac{\omega - \omega_k}{kv \cos \theta}, \tag{22}$$

when the following inequality satisfied

$$\left| \frac{\omega - \omega_k}{kv \cos \theta} \right| < 1. \tag{23}$$

$$M = \frac{2}{\pi |kv \cos \theta \sin(\omega_0 t_0)|}$$

Using (22) and the last expression one can get the multiplier

$$M = \frac{2}{\pi \left| kv \cos \theta \sqrt{1 - \frac{(\omega - \omega_k)^2}{k^2 v^2 \cos^2 \theta}} \right|} = \frac{2}{\pi \sqrt{k^2 v^2 \cos^2 \theta - (\omega - \omega_k)^2}}. \tag{24}$$

Using this expression one can get

$$F(\omega)=\frac{2}{\pi}\int\limits_{0}^{+\infty}\int\limits_{0}^{2\pi}\frac{\Phi(k,\theta)d\theta kdk}{\sqrt{k^2v^2\cos^2\theta-(\omega-\omega(k))^2}}. \tag{25}$$

It is necessary to understand that the limits of integration have to be chosen in order to satisfy inequality (23).

4 Convenient set of dimensionless variables

It is convenient to introduce a new variable and a constant

$$\zeta=\frac{kv^2}{g},\ dk=\frac{g}{v^2}d\zeta,\ \lambda=\frac{v\omega}{g}. \tag{26}$$

One can see that ζ is a dimensionless analogue of k, and λ is just the same for ω. In these variables the spectrum Eq. (25) takes the following relatively simple form

$$F(\lambda)=\frac{2}{\pi}\frac{g}{v^3}B\int\limits_{0}^{+\infty}\int\limits_{0}^{2\pi}\frac{\Phi(\zeta,\theta)d\zeta d\theta}{\sqrt{\cos^2\theta-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}}, \tag{27}$$

where B is the constant which appears as a result of the substitution $k\to\zeta$ in $\Phi(k)$. For the Phillips spectrum case Eq. (17) this constant takes the form $B=\alpha v^8/g^4$.

The physical meaning of the constant λ becomes more obvious if we transform it to the following form

$$\lambda=\frac{\omega}{\omega_p}\frac{v}{c_p}, \tag{28}$$

where $\omega_p=g/c_p$ is the peak-frequency (this is the only frequency-like scale in the model) and c_p is the phase velocity. The ratio $\epsilon=v/c_p$ according to Banner (1990) is 0.05 in magnitude.

For the case of strongly anisotropic spectrum ($\Phi(k,\theta)=\Phi(k)\delta(\theta)$) one can easily obtain the following result by a simple substitution in Eq. (27)

$$F(\lambda)=\frac{2}{\pi}\frac{g}{v^3}B\int\limits_{0}^{+\infty}\frac{\Phi(\zeta)d\zeta}{\sqrt{1-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}}. \tag{29}$$

If the spectrum is isotropic ($\Phi(k,\theta)=\Phi(k)$) things become a little more complex. Following Eq. (27) we have

$$F(\lambda)=\frac{8}{\pi}\frac{g}{v^3}B\int\limits_{0}^{+\infty}\Phi(\zeta)d\zeta\int\limits_{0}^{\pi/2}\frac{d\theta}{\sqrt{\cos^2\theta-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}}, \tag{30}$$

An analysis of the θ-part gives

$$\int\limits_0^{\pi/2} \frac{d\theta}{\sqrt{\cos^2\theta - \left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}} =$$

$$= \frac{1}{\sqrt{1-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}} K\left(\frac{1}{\sqrt{1-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}}\right) =$$

$$= K\left(\sqrt{1-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}\right) + iK\left(\frac{\lambda-\lambda_k}{\zeta}\right), \tag{31}$$

where K is the complete elliptic integral of the first kind. Using the inequality (23) after substitution Eq. (31) into Eq. (30) one can obtain

$$F(\lambda) = \frac{8}{\pi}\frac{g}{v^3}B\int\limits_0^{+\infty}\Phi(\zeta)d\zeta K\left(\sqrt{1-\left(\frac{\lambda-\lambda_k}{\zeta}\right)^2}\right). \tag{32}$$

Expressions (29) and (32) are valid for an arbitrary dispersion relation $\lambda_k = \omega(k)v/g$.

5 Deep water case

For further study of the isotropic spectrum (this case is more complex so let us do it first and then perform the same calculations for the anisotropic case) we have to introduce the form of the dispersion law. We think that the case of the deep water surface is the most important. We have to use explicit expression of ω_k, in the case of a deep water $\omega_k = \sqrt{gk}$ or $\lambda_k = \sqrt{\zeta}$

$$F(\lambda) = \frac{8}{\pi}\frac{g}{v^3}B\int\limits_0^{+\infty}\Phi(\zeta)d\zeta K\left(\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}\right). \tag{33}$$

Now we have to take into account condition (23)

$$\zeta^2 - \left(\lambda - \sqrt{\zeta}\right)^2 \geq 0. \tag{34}$$

Solving this inequality gives us

$$\begin{aligned} y_{1,2} &= \frac{1}{2} \pm \sqrt{\frac{1}{4} - \lambda}, \\ y_{3,4} &= -\frac{1}{2} \pm \sqrt{\frac{1}{4} + \lambda}, \end{aligned} \tag{35}$$

where $y = \sqrt{\zeta}, y \geq 0$, so we can state that interesting domains are

$$\text{when} \;\; \lambda \leq \frac{1}{4},$$

$$F(\lambda) = \frac{8}{\pi}\frac{g}{v^3}B\left[\int\limits_{\left(-\frac{1}{2}+\sqrt{\frac{1}{4}+\lambda}\right)^2}^{\left(\frac{1}{2}-\sqrt{\frac{1}{4}-\lambda}\right)^2} \Phi(\zeta)K\left(\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}\right)d\zeta \; + \right.$$
$$\left. + \int\limits_{\left(\frac{1}{2}+\sqrt{\frac{1}{4}-\lambda}\right)^2}^{+\infty} \Phi(\zeta)K\left(\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}\right)d\zeta\right]; \quad (36)$$

$$\text{when} \;\; \lambda > \frac{1}{4},$$

$$F(\lambda) = \frac{8}{\pi}\frac{g}{v^3}B \int\limits_{\left(-\frac{1}{2}+\sqrt{\frac{1}{4}+\lambda}\right)^2}^{+\infty} \Phi(\zeta)d\zeta K\left(\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}\right). \quad (37)$$

Let us explore special case, when $\Phi(\zeta)$ is a quickly going down to zero function, like Phillips spectrum $\Phi(\zeta) = 1/\zeta^4$. In this case we can take into account only first part of (36) at $\lambda \ll 1/4$. Calculation of the spectrum decomposition is situated in Appendix A and (A7) gives us the following result

$$F(\lambda) = F_0(1 + \frac{1}{4}\lambda^2 + ...), \quad (38)$$

where

$$F_0 = 8\pi\alpha\frac{v^5}{g^3}\frac{1}{\lambda^5}, \quad (39)$$

this is just the Phillips spectrum Eq. (18) in absence of Doppler effect for isotropic distribution.

For the one-directional spectra we can use exactly the same approach, but all calculations become simpler. The starting domains of integrations (which correspond to Eq. (36) and Eq. (37)) are the following

$$\text{when} \;\; \lambda \leq \frac{1}{4},$$

$$F(\lambda) = \frac{2}{\pi}\frac{g}{v^3}B\left(\int\limits_{\left(-\frac{1}{2}+\sqrt{\frac{1}{4}+\lambda}\right)^2}^{\left(\frac{1}{2}-\sqrt{\frac{1}{4}-\lambda}\right)^2} \Phi(\zeta)\frac{d\zeta}{\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}} + \right.$$
$$\left. + \int\limits_{\left(\frac{1}{2}+\sqrt{\frac{1}{4}-\lambda}\right)^2}^{+\infty} \frac{\Phi(\zeta)d\zeta}{\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}}\right); \quad (40)$$

when $\lambda > \dfrac{1}{4}$,

$$F(\lambda) = \frac{2}{\pi}\frac{g}{v^3}B \int\limits_{\left(-\frac{1}{2}+\sqrt{\frac{1}{4}+\lambda}\right)^2}^{+\infty} \frac{\Phi(\zeta)d\zeta}{\sqrt{1-\left(\frac{\lambda-\sqrt{\zeta}}{\zeta}\right)^2}}. \tag{41}$$

The complete analysis of this case is situated in Appendix A and (A9) gives us the following formula

$$F(\lambda) = F_0(1 + \frac{1}{2}\lambda^2 + ...), \tag{42}$$

where

$$F_0 = 4\alpha\frac{v^5}{g^3}\frac{1}{\lambda^5}, \tag{43}$$

this is just the Phillips spectrum Eq. (19) in absence of Doppler effect for strongly anisotropic distribution.

Let us try to evaluate the range of applicability of the obtained results. As it was postulated in Eq. (28) we can state

$$\lambda = \frac{\omega}{\omega_0}\frac{v}{c_p} \leq \frac{1}{4}.$$

Following Banner (1990) we can take $\epsilon = v/c_p = 0.05$. Using this one can obtain the following inequality

$$\omega \leq 5\omega_0. \tag{44}$$

It means that we cannot consider the ω^{-4} spectrum as an artifact of the observation of time histories at a point brought about by Doppler shifting of the short waves riding on the orbital currents of the long waves.

6 Numerical results

The integrals Eq. (36), Eq. (37), Eq. (40) and Eq. (41) can be easily calculated numerically. It is not very illustrative to view the explicit form of the spectrum, so let us represent it as the ratio $F(\lambda)/F_0(\lambda)$. The expressions Eq. (36), Eq. (40) and Eq. (37), Eq. (41) give us the curves shown in Fig. 1.

It is obvious that the form of the first part of the decomposition is not linear and it means that we cannot transform the ω^{-5} to ω^{-4}. The most illustrative form to estimate the power of the ω in $F(\lambda)$ is the logarithm scale. The results are represented in Fig. 2

As one can easily see, there is almost no difference betwin ω^{-5} line and calculated curve. Thus in this range ($\lambda \leq 1/4$) we can state that the does not give Doppler a desirable result. We can see that there are no prerequisites for transformation Phillips spectrum (ω^{-5}) to experimentally observed ω^{-4}.

7 Conclusion

The presence of at least two different approaches on the wind-driven sea waves frequency spectra formation motivates us to examine the Phillips model in the presence of the Doppler effect. The pure Phillips model gives the results essentially different with the experimental data. Following Banner conjecture we obtained the general formula for an arbitrary frequency spectrum in the presence of the Doppler effect. Using this formula we have performed an analytical analysis of the influence of the Doppler frequency shifting on surface waves' spectra at the deep water and strongly anisotropic cases. The forms of spectra were calculated numerically for a wide range of frequencies.

Analytical and numerical results give us an irrefutable proof of inconsistency of Doppler shifting mechanism in fitting Phillips model with the experimental results. An analytical estimation shows that in the presence of Doppler effect its contribution is negligible. The numerical computation gives spectrum decay $\sim \omega^{-5}$ at least up to frequency $\sim 20\omega_p$.

According to the present experimental results one has to recognize that Phillips model of energy dissipation does not give the result similar to experimental data even if the Doppler effect takes place.

Appendix A Calculations of the spectra decompositions

At first, a little bit more complex case of isotropic spectrum. Let us perform the following substitution

$$x = \frac{\lambda - \sqrt{\zeta}}{\zeta}. \tag{A1}$$

In this case in our domain of integration ($-1 < x < +1$) we can obtain the invert representation

$$\zeta = \frac{1}{2x^2}\left[1 - 2\sqrt{1+4x\lambda} + 2x\lambda\right], \tag{A2}$$

and further

$$\zeta(x) \approx \frac{1}{2x^2}\left[2\lambda^2x^2 - 4\lambda^3x^3 + 10\lambda^4x^4\right] = \lambda^2 - 2\lambda^3x + 5\lambda^4x^2. \tag{A3}$$

For substitution $d\zeta = \zeta'(x)dx$ we have to calculate the first derivative

$$\begin{aligned}
\zeta'(x) = -\frac{1}{x^3} - \frac{\lambda}{x^2\sqrt{1+4\lambda x}} + \frac{\sqrt{1+4\lambda x}}{x^3} - \\
- \frac{\lambda}{x^2} \approx -\frac{1}{x^3} - \frac{\lambda}{x^2}\left[1 - 2\lambda x + 6\lambda^2x^2 - 20\lambda^3x^3 + \right. \\
\left. + 70\lambda^4x^4\right] + + \frac{1}{x^3}\left[1 + 2\lambda x - 2\lambda^2x^2 + 4\lambda^3x^3 - \right. \\
\left. - 10\lambda^4x^4 + 28\lambda^5x^5\right] - \frac{\lambda}{x^2} = \\
= -2\lambda^3\left(1 - 5\lambda x + 21\lambda^2x^2\right).
\end{aligned} \tag{A4}$$

$$\begin{aligned}
\Phi(\zeta(x)) = \Phi(\lambda^2(1 - 2\lambda x + 5\lambda^2x^2)) \approx \\
\approx \Phi(\lambda^2) - \frac{\Phi'(\lambda^2)}{1!}2\lambda^3x + \\
+ \frac{\Phi'(\lambda^2)}{1!}5\lambda^4x^2 + \frac{\Phi''(\lambda^2)}{2!}4\lambda^6x^2.
\end{aligned} \tag{A5}$$

Now we can calculate the integral (36) with Phillips spectrum ($\Phi(\zeta) = 1/\zeta^4$)

$$F(\lambda) = \frac{8}{\pi}\frac{g}{v^3}B2\lambda^3 \int\limits_{-1}^{+1} \Phi(\zeta(x))\zeta'(x)K(\sqrt{1-x^2})dx, \tag{A6}$$

Taking into account several well known results

$$\int\limits_{-1}^{+1} K\left(\sqrt{1-x^2}\right) dx = \frac{\pi^2}{2},$$

$$\int\limits_{-1}^{+1} xK\left(\sqrt{1-x^2}\right) dx \equiv 0,$$

$$\int\limits_{-1}^{+1} x^2K\left(\sqrt{1-x^2}\right) dx = \int\limits_{0}^{+1} E\left(\sqrt{1-x^2}\right) dx = \frac{\pi^2}{8},$$

where E is the complete elliptic integral of the second kind, one can obtain the final result

$$F(\lambda) = 8\pi\alpha\frac{v^5}{g^3}\frac{1}{\lambda^5}(1 + \frac{1}{4}\lambda^2 + ...). \tag{A7}$$

Calculations of the spectrum decomposition for strongly anisotropic case up to expression Eq. (A6) are just the same. Integral (40) takes the form

$$F(\lambda) = \frac{2}{\pi}\frac{g}{v^3}B2\lambda^3 \int\limits_{-1}^{+1} \frac{\Phi(\zeta(x))\zeta'(x)}{\sqrt{1-x^2}}dx, \tag{A8}$$

The final result is

$$F(\lambda) = 2\alpha\frac{v^5}{g^3}\frac{1}{\lambda^5}(1 + \frac{1}{2}\lambda^2 + ...). \tag{A9}$$

Acknowledgements. The author gratefully wishes to acknowledge the following contributions: set up of this problem and many enlightening and helpful discussions by Prof. V.E. Zakharov; Landau Scholarship Committee; grant INTAS-00-00292; grant RFBR 00-15-96007.

References

Banner, M.L., Equilibrium Spectra of Wind Generated Waves. Journ. of Physical Oceanography, 20, 966-984,1990

Craik, A.D.D., Interaction of a short-wave field with a dominant long wave in deep water: derivation from Zakharov's spectral formulation. Journ. Austral. Math. Soc. Ser. B, 29, 430-439,1987

Donelan, M.A., Hamilton J. and Hui W.H., Directional spectra of wind generated waves. Phil. Trans. R. Soc. London, A315, 509-562, 1985

Donelan, M.A., Air-water exchange processes. In: Physical Processes in Oceans and Lakes (J. Imberger, Ed.), AGU Coastal and Estuarine Studies, 54, 19-36, 1998

Forristall, G.Z., Measurements of a saturated range in ocean wave spectra. J. Geophys. Res., 86, c9, 8075-8084, 1981

Hansen at al., J. Phys. Oceanogr., 20, 1264-1277.

Hasselman, K., On the nonlinear energy transfer in a gravity wave spectrum. Part 1. J. Fluid Mech., 12, 481-500, 1962

Hasselman, K., On the nonlinear energy transfer in a gravity wave spectrum. Part 2 and Part 3. J. Fluid Mech., 15, 273-281 and 385-398,1963

Kahma, K.K., A study of the growth of the wave spectrum with fetch. J. Phys. Oceanogr., 11, 1503-1515, 1981

Phillips, O.M., The equilibrium range in the spectrum of wind generated waves. J. Fluid Mech., 4, 426-434, 1958

Pushkarev, A.N. and Zakharov, V.E., Phys. Rev. Lett., 76, 1996

Pushkarev, A.N. and Zakharov, V.E., Physica D, 145, 2000

Toba, Y., Local balance in the air-sea boundary processes Part II: On the spectrum of wind generated waves. J. Oceanogr. Soc. Japan, 29, 209-220, 1973

Zakharov, V.E. and Filonenko, N.N., Dokl. Acad. Nauk SSSR, 170, 6, 1966 881-883,1968

Zakharov, V.E. and Filonenko, N.N., J. Appl. Mech. Tech. Phys, 4, 1967

Zakharov, V.E. and Filonenko, N.N., Energy spectrum for stochastic oscillations of the surface of a liquid. Soviet Phys. Dokl., 11, 881-883,1968

Zakharov, V.E., Falkovich, G., Lvov, V., Kolmogorov spectra of turbulence I (Springer, Berlin), 1992

Figure Captions

Fig. 1. Ratio of calculated spectra F to Phillips spectrum as a function of dimensionless variable λ is shown. It is clearly that F starting to deviate from Phillips spectrum only at $\lambda \sim 0.4$

Fig. 2. Logarithm of calculated spectra F and Phillips spectrum as a function of logarithm of dimensionless variable λ is represented. It is easy to see that calculated spectra are lines are exactly parallel to line corresponding to Phillips spectra. This allow us to conclude that the power of frequency is the same and is equal to -5

Figures



Fig. 1. Ratio of calculated spectra F to Phillips spectrum as a function of dimensionless variable λ is shown. It is clearly that F starting to deviate from Phillips spectrum only at $\lambda \sim 0.4$



Fig. 2. Logarithm of calculated spectra F and Phillips spectrum as a function of logarithm of dimensionless variable λ is represented. It is easy to see that calculated spectra are lines are exactly parallel to line corresponding to Phillips spectra. This allow us to conclude that the power of frequency is the same and is equal to -5